

Question #1 of 82

Question ID: 438835

The table below shows a sample of returns on two securities:

Period	1	2	3	4	Mean
Security P	0.2%	0.5%	1.1%	-0.6%	0.3%
Security Q	-0.3%	0.9%	1.5%	-0.5%	0.4%

The sample covariance between the two securities' returns is closest to:

- ☐ A) 0.78.
- ☐ B) 1.86.
- ☐ C) 0.47.
- ☒ D) 0.62.

Explanation

Period	1	2	3	4	Sum
$P_i - \bar{P}$	-0.1	0.2	0.8	-0.9	
$Q_i - \bar{Q}$	-0.7	0.5	1.1	-0.9	
$(P_i - \bar{P})(Q_i - \bar{Q})$	0.07	0.10	0.88	0.81	1.86

Sample covariance = $\frac{\sum_{i=1}^n (P_i - \bar{P})(Q_i - \bar{Q})}{n - 1} = \frac{1.86}{3} = 0.62$

Question #2 of 82

Question ID: 438853

Robert Patterson, an options trader, believes that the return on options trading is higher on Mondays than on other days. In order to test his theory, he formulates a null hypothesis. Which of the following would be an appropriate null hypothesis? Returns on Mondays are:

- ☐ A) not less than returns on other days.
- ☐ B) less than returns on other days.
- ☒ C) not greater than returns on other days.
- ☐ D) greater than returns on other days.

Explanation

An appropriate null hypothesis is one that the researcher wants to reject. If Patterson believes that the returns on Mondays are greater than on other days, he would like to reject the hypothesis that the opposite is true-that returns on Mondays are not greater than returns on other days.

Question #3 of 82

Question ID: 438878

Which of the following statements about hypothesis testing is *most* accurate?

- ✓ **A) The probability of a Type I error is equal to the significance level of the test.**
- ✗ **B)** To test the claim that X is greater than zero, the null hypothesis would be $H_0: X > 0$.
- ✗ **C)** If you can disprove the null hypothesis, then you have proven the alternative hypothesis.
- ✗ **D)** The power of a test is one minus the probability of a Type I error.

Explanation

The probability of getting a test statistic outside the critical value(s) when the null is true is the level of significance and is the probability of a Type I error. The power of a test is 1 minus the probability of a Type II error. Hypothesis testing does not prove a hypothesis, we either reject the null or fail to reject it. The appropriate null would be " $X \leq 0$ " with " $X > 0$ " as the alternative hypothesis.

Questions #4-5 of 82

Margo Hinsdale is testing the null hypothesis that the population mean is less than or equal to 45. A random sample of 81 observations selected from this population produced a mean of 46.3. The population has a standard deviation of 4.5.

Question #4 of 82

Question ID: 438890

The value of the *appropriate* test statistic for the test of the population mean is:

- ✗ **A) $t = 4.60$.**
- ✗ **B) $z = -2.75$.**
- ✓ **C) $z = 2.60$.**
- ✗ **D) $t = 3.84$.**

Explanation

The population variance is known and the sample size is large. The test statistic is:

$$z = \frac{(46.3 - 45)}{4.5 / \sqrt{81}} = 2.60$$

Question #5 of 82

Question ID: 438891

At a 1 percent level of significance, the correct decision is to:

- ✓ **A) reject the null hypothesis.**
- ✗ **B)** neither reject nor fail to reject the null hypothesis.
- ✗ **C)** fail to reject the null hypothesis.
- ✗ **D)** accept the null hypothesis.

Explanation

Decision rule: reject H_0 if $z_{\text{computed}} > z_{\text{critical}}$. Therefore, reject the null hypothesis because the computed test statistic of 2.60 (see the answer to Part 1) exceeds the critical z -value of 2.33.

Questions #6-9 of 82

An analyst is testing to see if the mean of a population is less than 133. A random sample of 50 observations had a mean of 130. Assume a standard deviation of 5. The test is to be made at the 1% level of significance.

<i>z</i>	<i>0.00</i>	<i>0.01</i>	<i>0.02</i>	<i>0.03</i>	<i>0.04</i>	<i>0.05</i>	<i>0.06</i>
<i>0.0</i>	0.0000	0.0040	0.0080	0.0120	0.0160	0.0199	0.0239
<i>0.1</i>	0.0398	0.0438	0.0478	0.0517	0.0557	0.0596	0.0636
<i>0.2</i>	0.0793	0.0832	0.0871	0.0910	0.0948	0.0987	0.1026
<i>0.3</i>	0.1179	0.1217	0.1255	0.1293	0.1331	0.1368	0.1406
<i>1.7</i>	0.4554	0.4564	0.4573	0.4582	0.4591	0.4599	0.4608
<i>1.8</i>	0.4641	0.4649	0.4656	0.4664	0.4671	0.4678	0.4686
<i>1.9</i>	0.4713	0.4719	0.4726	0.4732	0.4738	0.4744	0.4750
<i>2.0</i>	0.4772	0.4778	0.4783	0.4788	0.4793	0.4798	0.4803
<i>2.1</i>	0.4821	0.4826	0.4830	0.4834	0.4838	0.4842	0.4846
<i>2.2</i>	0.4861	0.4864	0.4868	0.4871	0.4875	0.4878	0.4881
<i>2.3</i>	0.4893	0.4896	0.4898	0.4901	0.4904	0.4906	0.4909
<i>2.4</i>	0.4918	0.4920	0.4922	0.4925	0.4927	0.4929	0.4931

Question #6 of 82

Question ID: 438861

The null hypothesis is:

- X A) $\mu > 133$.
- ✓ B) $\mu \geq 133$.
- X C) $\mu \leq 133$.
- X D) $\mu = 133$.

Explanation

The null hypothesis is the hypothesis that the researcher wants to reject. Here the hypothesis that is being looked for is that the mean of a population is less than 133. The null hypothesis is that the mean is greater than or equal to 133. The question is whether the null hypothesis will be rejected in favor of the alternative hypothesis that the mean is less than 133.

Question #7 of 82

Question ID: 438862

The calculated test statistic is:

- X A) +1.33.
- X B) -3.00.
- X C) -1.33.
- ✓ D) -4.24.

Explanation

A test statistic is calculated by subtracting the hypothesized parameter from the parameter that has been estimated and dividing

the difference by the standard error of the sample statistic. Here, the test statistic = (sample mean - hypothesized mean) / ((sample standard deviation / (sample size)^{1/2})) = (130 - 133) / (5 / 50^{1/2}) = (-3) / (5 / 7.0711) = -4.24.

Question #8 of 82

Question ID: 438863

The critical value is:

- ☐ A) -2.38.
- ☐ B) 2.47.
- ☒ C) -2.33.
- ☐ D) 2.17.

Explanation

This is a one-tailed test with a significance level of 0.01. The critical value for a one-tailed test at a 1% level of significance is -2.33.

Question #9 of 82

Question ID: 438864

You should:

- ☐ A) Cannot be determined with the information given.
- ☐ B) accept the null hypothesis.
- ☐ C) reject the alternative hypothesis.
- ☒ D) reject the null hypothesis.

Explanation

The calculated test statistic of -4.24 falls to the left of the z-statistic of -2.33, and is in the rejection region. Thus, the null hypothesis is rejected and the conclusion is that the population mean is less than 133.

Question #10 of 82

Question ID: 438852

Brian Ci believes that the average return on equity in the airline industry, μ , is less than 5%. What are the appropriate null (H_0) and alternative (H_a) hypotheses to test this belief?

- ☐ A) $H_0: \mu > 0.05$ versus $H_a: \mu < 0.05$.
- ☐ B) $H_0: \mu < 0.05$ versus $H_a: \mu > 0.05$.
- ☒ C) $H_0: \mu \geq 0.05$ versus $H_a: \mu < 0.05$.
- ☐ D) $H_0: \mu < 0.05$ versus $H_a: \mu \geq 0.05$.

Explanation

The alternative hypothesis is determined by the theory or the belief. The researcher specifies the null as the hypothesis that he wishes to reject (in favor of the alternative). Note that this is a one-sided alternative because of the "less than" belief.

Question #11 of 82

Question ID: 438832

A sample covariance of two random variables is most commonly utilized to:

- ☐ **A) demonstrate either the presence or absence of spurious correlation of the variables.**
- ☐ **B) estimate the "pure" measure of the tendency of two variables to move together over a period of time.**
- ☐ **C) identify and measure strong nonlinear relationships between the two variables.**
- ☒ **D) calculate the correlation coefficient, which is a measure of the strength of their linear relationship.**

Explanation

Since the actual value of a sample covariance can range from negative to positive infinity depending on the scale of the two variables, it is most commonly used to calculate a more useful measure, the correlation coefficient.

Question #12 of 82

Question ID: 438898

The test statistic for an F-test of the equality of two sample variances is the:

- ☐ **A) product of the two sample standard deviations.**
- ☒ **B) ratio of the two sample variances.**
- ☐ **C) product of the two sample variances.**
- ☐ **D) ratio of the two sample standard deviations.**

Explanation

The test statistic for an F-test of the equality of two sample variances is the ratio of the two sample variances.

$$F = s_1^2 / s_2^2.$$

Question #13 of 82

Question ID: 438849

The approximate 99% confidence interval for the population mean based on a sample of 60 returns with a mean of 7% and a sample standard deviation of 25% is *closest* to:

- ☐ **A) 1.546% to 13.454%.**
- ☐ **B) 0.546% to 13.454%.**
- ☒ **C) -1.584% to 15.584%.**
- ☐ **D) 1.584% to 14.584%.**

Explanation

The standard error for the mean = $s / (n)^{0.5} = 25\% / (60)^{0.5} = 3.227\%$. The critical value from the t-table should be based on $60 - 1 = 59$ df. Since the standard tables do not provide the critical value for 59 df the closest available value is for 60 df. This leaves us with an approximate confidence interval. Based on 99% confidence and $df = 60$, the critical t-value is 2.660. Therefore the 99% confidence interval is approximately: $7\% \pm 2.660(3.227)$ or $7\% \pm 8.584\%$ or -1.584% to 15.584% .

If you use a z-statistic, the confidence interval is $7\% \pm 2.58(3.227) = -1.326\%$ to 15.326% , which is closest to the correct choice.

Questions #14-18 of 82

An analyst is conducting a hypothesis test to determine if the mean time spent on investment research is different from 3 hours

per day. The test is performed at the 5 percent level of significance and uses a random sample of 64 portfolio managers, where the mean time spent on research is found to be 2.5 hours. The population standard deviation is 1.5 hours.

Question #14 of 82

Question ID: 438884

The appropriate null hypothesis for the described test is:

- ☐ A) $H_0: \mu \geq 3$ hours.
- ☐ B) $H_0: \mu \neq 3$ hours.
- ☒ C) $H_0: \mu = 3$ hours.
- ☐ D) $H_0: \mu \leq 3$ hours.

Explanation

$H_0: \mu = 3$ hours.

Question #15 of 82

Question ID: 438885

Which type of test is this?

- ☒ A) Two-tailed test.
- ☐ B) Chi-square test.
- ☐ C) One-tailed test.
- ☐ D) Equity of variance test.

Explanation

This is a two-sided (tailed) test. We want to test if the mean "differs from" 3 hours (i.e., $H_a: \mu \neq 3$ hours).

Question #16 of 82

Question ID: 438886

The calculated z-statistic is:

- ☐ A) +2.67.
- ☐ B) -2.13.
- ☐ C) +0.33.
- ☒ D) -2.67.

Explanation

The normally distributed test statistic is:

Question #17 of 82

Question ID: 438887

The critical z-value(s) of the test statistic is (are):

- ☒ A) ± 1.96 .
- ☐ B) -1.96.
- ☐ C) ± 2.58 .
- ☐ D) +1.96.

Explanation

At $\alpha/2 = 0.025$, the critical z-values are: $\pm z_{\alpha/2} = \pm z_{0.025} = \pm 1.96$.

Question #18 of 82

Question ID: 438888

Which of the following decisions is the **CORRECT** decision for this study?

- ☐ A) Fail to reject the null hypothesis.
- ☒ B) Reject the null hypothesis.
- ☐ C) No decision is possible because the sample standard deviation was not given.
- ☐ D) The sample size is too small, so increase the sample size.

Explanation

Decision rule: reject H_0 if $z_{\text{computed}} < -1.96$, or $z_{\text{computed}} > +1.96$. Since $-2.67 < -1.96$, reject H_0 .

Question #19 of 82

Question ID: 438899

What is the computed value of the test statistic that follows an F -distribution when sample variances are equal and the level of significance is 0.10?

- ☐ A) 0.90.
- ☐ B) 0.05.
- ☒ C) 1.00.
- ☐ D) 0.10.

Explanation

The calculated value of the F -distributed test statistic, F , is s_1^2 / s_2^2 . With equal variances, $F = 1$.

Question #20 of 82

Question ID: 438846

The average U.S. dollar/Euro exchange rate from a sample of 36 monthly observations is \$1.00/Euro. The population variance is 0.49. What is the 95% confidence interval for the mean U.S. dollar/Euro exchange rate?

- ☐ A) \$0.8657 to \$1.1343.
- ☐ B) \$0.5100 to \$1.4900.
- ☒ C) \$0.7713 to \$1.2287.
- ☐ D) \$0.8075 to \$1.1925.

Explanation

The population *standard deviation* is the square root of the variance ($\sqrt{0.49} = 0.7$). Because we know the population standard deviation, we use the z-statistic. The z-statistic reliability factor for a 95% confidence interval is 1.960. The confidence interval is $\$1.00 \pm 1.960(\$0.7 / \sqrt{36})$ or $\$1.00 \pm \$0.2287$.

Question #21 of 82

Question ID: 438834

You observe the following pairs of annual returns on a sample of two stocks: [(0.03, 0.01), (0.06, 0.06), (0.02, 0.05), (0.13,

0.08)]. The covariance of the returns on the two stocks is *closest* to:

- ✓ **A) 0.0011.**
- X **B) 0.0033.**
- X **C) 0.00165.**
- X **D) 0.0008.**

Explanation

The means for the two stocks are:

$$(0.03 + 0.06 + 0.02 + 0.13) / 4 = 0.06 \text{ and } (0.01 + 0.06 + 0.05 + 0.08) / 4 = 0.05.$$

The covariance is:

$$[(0.03 - 0.06)(0.01 - 0.05) + (0.06 - 0.06)(0.06 - 0.05) + (0.02 - 0.06)(0.05 - 0.05) + (0.13 - 0.06)(0.08 - 0.05)] / [4 - 1] = 0.0011.$$

Question #22 of 82

Question ID: 438868

A Type I error occurs when the null hypothesis:

- X **A) fails to be rejected when it is false.**
- ✓ **B) is rejected when it is true.**
- X **C) is rejected when it is false.**
- X **D) fails to be rejected when it is true.**

Explanation

A Type I error occurs when the null hypothesis is rejected when it is true. A Type II error occurs when the null hypothesis fails to be rejected when it is false.

Question #23 of 82

Question ID: 438859

An analyst conducts a two-tailed z-test to determine if small cap returns are significantly different from 10%. The sample size was 200. The computed z-statistic is 2.3. Using a 5% level of significance, which statement is *most* accurate?

- X **A) Fail to reject the null hypothesis and conclude that small cap returns are close enough to 10% that we cannot say they are significantly different from 10%.**
- X **B) You cannot determine what to do with the information given.**
- ✓ **C) Reject the null hypothesis and conclude that small cap returns are significantly different from 10%.**
- X **D) A sample size of 200 indicates that we should fail to reject the null.**

Explanation

At the 5% level of significance the critical z-statistic for a two-tailed test is 1.96 (assuming a large sample size). The null hypothesis is $H_0: x = 10\%$. The alternative hypothesis is $H_A: x \neq 10\%$. Because the computed z-statistic is greater than the critical z-statistic ($2.33 > 1.96$), we reject the null hypothesis and we conclude that small cap returns are significantly different than 10%.

Question #24 of 82

Question ID: 438876

The mean monthly return for an equity portfolio over 60 months is 1.5 percent. The standard deviation is 3.0 percent. The value of the test statistic to test the hypothesis that mean monthly return is equal to zero is *closest* to:

- ✓ **A) 3.87.**
- X **B) 30.00.**
- X **C) 2.19.**
- X **D) 0.50.**

Explanation

$$z = (1.5\% - 0.0\%) / [(3.0\% / \sqrt{60})] = 3.87$$

Question #25 of 82

Question ID: 438873

A survey is taken to determine whether the average starting salaries of CFA charterholders is equal to or greater than \$59,000 per year. What is the test statistic given a sample of 135 newly acquired CFA charterholders with a mean starting salary of \$64,000 and a standard deviation of \$5,500?

- X **A) -0.91.**
- X **B) -10.56.**
- ✓ **C) 10.56.**
- X **D) 0.91.**

Explanation

With a large sample size (135) the z-statistic is used. The z-statistic is calculated by subtracting the hypothesized parameter from the parameter that has been estimated and dividing the difference by the standard error of the sample statistic. Here, the test statistic = (sample mean - hypothesized mean) / (population standard deviation / (sample size)^{1/2}) = $(\bar{X} - \mu) / (\sigma / n^{1/2}) = (64,000 - 59,000) / (5,500 / 135^{1/2}) = (5,000) / (5,500 / 11.62) = 10.56$.

Question #26 of 82

Question ID: 438882

Which of the following statements about hypothesis testing is *most* accurate?

- X **A) When the critical Z-statistic is greater than the sample Z-statistic in a two-tailed test, reject the null hypothesis and accept the alternative hypothesis.**
- X **B) A Type I error is rejecting the null hypothesis when it is true, and a Type II error is accepting the alternative hypothesis when it is false.**
- ✓ **C) A hypothesized mean of 3, a sample mean of 6, and a standard error of the sampling means of 2 give a sample Z-statistic of 1.5.**
- X **D) A two-tailed test on a large sample with a significance level of 0.01 has confidence intervals of ± 1.96 standard errors.**

Explanation

The 0.01 level of significance has confidence intervals of ± 2.58 standard errors. A Type II error is wrongly accepting the null hypothesis. The null hypothesis should be rejected when the sample Z-statistic is greater than the critical Z-statistic.

Question #27 of 82

Question ID: 438851

Susan Bellows is comparing the return on equity for two industries. She is convinced that the return on equity for the discount retail industry (DR) is greater than that of the luxury retail (LR) industry. What are the hypotheses for a test of her comparison of return on equity?

- ✓ **A) $H_0: \mu_{DR} \leq \mu_{LR}$ versus $H_a: \mu_{DR} > \mu_{LR}$.**
- X **B) $H_0: \mu_{DR} = \mu_{LR}$ versus $H_a: \mu_{DR} \neq \mu_{LR}$.**
- X **C) $H_0: \mu_{DR} \neq \mu_{LR}$ versus $H_a: \mu_{DR} = \mu_{LR}$.**
- X **D) $H_0: \mu_{DR} = \mu_{LR}$ versus $H_a: \mu_{DR} < \mu_{LR}$.**

Explanation

The alternative hypothesis is determined by the theory or the belief. The researcher specifies the null as the hypothesis that she wishes to reject (in favor of the alternative). Note that this is a one-sided alternative because of the "greater than" belief.

Questions #28-32 of 82

Austin Roberts wants to test if the mean price of houses in the area is greater than \$145,000. A random sample of 36 houses in the area has a mean price of \$149,750. The population standard deviation is \$24,000, and Roberts wants to conduct hypothesis testing at a 1 percent level of significance.

Question #28 of 82

Question ID: 438893

The appropriate alternative hypothesis is:

- X **A) $H_a: \mu \pm \$145,000$.**
- ✓ **B) $H_a: \mu > \$145,000$.**
- X **C) $H_a: \mu < \$145,000$.**
- X **D) $H_a: \mu \geq \$145,000$.**

Explanation

$H_a: \mu > \$145,000$

Question #29 of 82

Question ID: 438894

The value of the calculated test statistic is *closest* to:

- X **A) $z = 0.67$.**
- X **B) $z = 8.13$.**
- X **C) $z = 4.00$.**
- ✓ **D) $z = 1.19$.**

Explanation

$$z = \frac{149,750 - 145,000}{24,000 / \sqrt{36}} = 1.1875$$

Question #30 of 82

Question ID: 438895

Which of the following *most accurately* describes the appropriate test structure?

- ✓ **A) One-tailed test.**
- X B) Two-tailed test.
- X C) F-test.
- X D) Chi-square test.

Explanation

The alternative hypothesis, $H_a: \mu > \$145,000$, only allows for values greater than the hypothesized value. Thus, this is a one-sided (one-tailed) test.

Question #31 of 82

Question ID: 438896

The critical value of the z-statistic is:

- X A) $z = \pm 2.33$.
- X B) $z = \pm 1.96$.
- ✓ C) $z = +2.33$.
- X D) $z = -2.33$.

Explanation

For a one-tailed z-test at the 1% level of significance, the critical z-value is $z_{0.01} = 2.33$. Since the test is one-tailed on the upper end (i.e., $H_a: \mu > 145,000$), we use a positive z-critical value

Question #32 of 82

Question ID: 438897

At a 1 percent level of significance, Roberts should:

- ✓ **A) fail to reject the null hypothesis.**
- X B) reject the null hypothesis.
- X C) accept the null hypothesis.
- X D) neither reject nor fail to reject the null hypothesis.

Explanation

The decision rule is: reject H_0 if $z\text{-computed} > z\text{-critical}$. Since $1.1875 < 2.33$, Roberts will fail to reject the null.

Question #33 of 82

Question ID: 438821

Assume that the following returns are a sample of annual returns for firms in the clothing industry. Given the following sample of returns, what are the sample variance and standard deviation?

Firm 1	Firm 2	Firm 3	Firm 4	Firm 5
15%	2%	5%	(7%)	0%

	<u>Variance</u>	<u>Standard Deviation</u>
✓ A) 64.5		8.0
X B) 51.6		7.2
X C) 22.0		4.7
X D) 32.4		5.7

Explanation

The sample variance is found by taking the sum of all squared deviations from the mean and dividing by $(n - 1)$. $[(15 - 3)^2 + (2 - 3)^2 + (5 - 3)^2 + (-7 - 3)^2 + (0 - 3)^2] / (5 - 1) = 64.5$

The sample standard deviation is found by taking the square root of the sample variance. $\sqrt{64.5} = 8.03$

Question #34 of 82

Question ID: 438870

Which of the following statements about testing a hypothesis using a Z-test is *least* accurate?

- X **A) If the calculated Z-statistic lies outside the critical Z-statistic range, the null hypothesis can be rejected.**
- X **B) A Type I error is rejecting the null hypothesis when it is actually true.**
- ✓ **C) The calculated Z-statistic determines the appropriate significance level to use.**
- X **D) The confidence interval for a two-tailed test of a population mean at the 5% level of significance is that the sample mean falls between $\pm 1.96 \sigma/\sqrt{n}$ of the null hypothesis value.**

Explanation

The significance level is chosen before the test so the calculated Z-statistic can be compared to an appropriate critical value.

Questions #35-37 of 82

John Starwall, CFA, is assigned the task of estimating the risk and return for security X, which is being considered as an addition to his current portfolio. If the investment is accepted, then the total portfolio will consist of five different equities that are not highly correlated. Security X has an estimated beta of 1.4. The probabilities of various states of the economy and the expected return for security X given each state are as follows:

<i>State of Economy</i>	<i>P_i</i>	<i>E(R_i)</i>
Recession	0.3	0.01
Average Economy	0.6	0.10
Boom	0.1	0.20

Question #35 of 82

Question ID: 438823

Based on the information above, what is the estimated expected return for security X?

- ☐ A) 15.5%.
- ☒ B) 8.3%.
- ☐ C) 10.0%.
- ☐ D) 8.0%.

Explanation

$$E(R) = 0.3(0.01) + 0.6(0.1) + 0.1(0.2) = 0.003 + 0.06 + 0.02 = 0.083 \text{ or } 8.3\%$$

Question #36 of 82

Question ID: 438824

Based on the information above, what is the estimated standard deviation for security X?

- ☐ A) 10.1%.
- ☒ B) 5.6%.
- ☐ C) 7.8%.
- ☐ D) 0.3%.

Explanation

$$\text{Variance} = 0.3(0.01 - 0.083)^2 + 0.6(0.10 - 0.083)^2 + 0.1(0.20 - 0.083)^2 = 0.00160 + 0.00017 + 0.00137 = 0.00314$$

$$\text{Standard Deviation} = (0.00314)^{0.5} = 0.056 \text{ or } 5.6\%$$

Question #37 of 82

Question ID: 438825

Based on the information above, what are the appropriate measures for assessing whether or not this security should be added to the portfolio?

- ☒ A) Expected return and standard deviation of the security.
- ☐ B) Beta and standard deviation of the security.
- ☐ C) Expected return, beta, and standard deviation of the security.
- ☐ D) Expected return and beta of the security.

Explanation

Total risk is comprised of systematic and unsystematic risk. In a well-diversified portfolio only systematic risk is important. Systematic risk is measured by beta. However, if the portfolio is not well diversified, total risk as measured by standard deviation must be considered. The stand-alone risk for this security is important in the decision because not all of the unsystematic risk is diversified away. Therefore, the correct answer is expected return and standard deviation. Beta only applies to a well-diversified portfolio.

Question #38 of 82

Question ID: 438858

An analyst conducts a two-tailed test to determine if mean earnings estimates are significantly different from reported earnings. The sample size is greater than 25 and the computed test statistic is 1.25. Using a 5% significance level, which of the following statements is *most* accurate?

- ☐ A) To test the null hypothesis, the analyst must determine the exact sample size and calculate the degrees of freedom for the test.

- X **B)** The analyst should reject the null hypothesis and conclude that the earnings estimates are significantly different from reported earnings.
- ✓ **C)** The analyst should fail to reject the null hypothesis and conclude that the earnings estimates are not significantly different from reported earnings.
- X **D)** The appropriate test to apply is a two-tailed chi-square test.

Explanation

The null hypothesis is that earnings estimates are equal to reported earnings. To reject the null hypothesis, the calculated test statistic must fall outside the two critical values. IF the analyst tests the null hypothesis with a z-statistic, the critical values at a 5% confidence level are ± 1.96 . Because the calculated test statistic, 1.25, lies between the two critical values, the analyst should fail to reject the null hypothesis and conclude that earnings estimates are not significantly different from reported earnings. If the analyst uses a t-statistic, the upper critical value will be even greater than 1.96, never less, so even without the exact degrees of freedom the analyst knows any t-test would fail to reject the null.

Questions #39-42 of 82

Two samples were drawn from a normally distributed population. For the first sample, the mean was \$50, and the standard deviation was \$5. For the second sample, the mean was \$55, and the standard deviation was \$6. The first sample consists of 25 observations, and the second sample consists of 36 observations. (Note: In the following questions, the subscripts "1" and "2" indicate the first and second sample, respectively.)

Question #39 of 82

Question ID: 438905

Consider the hypotheses structured as $H_0: \mu_1 = \$48$ versus $H_a: \mu_1 \neq \$48$. At a 1 percent level of significance, the null hypothesis:

- X **A) should be rejected.**
- X **B)** cannot be tested using this sample information provided.
- X **C)** should neither be rejected nor fail to be rejected.
- ✓ **D)** cannot be rejected.

Explanation

A two-tailed t-test is appropriate. The decision rule is to reject H_0 if the t-statistic is outside the range defined by $\pm t$ at $\alpha/2 = 0.025$ with $df = 24$. The t-statistic =

$$t_{24} = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} = \frac{50 - 48}{5/\sqrt{25}} = 2.0$$

$\pm t_{24}$ at $\alpha/2 = p = 0.025 = \pm 2.797$, therefore, H_0 cannot be rejected.

Question #40 of 82

Question ID: 438906

Using a 5 percent level of significance and a hypothesis test structure of $H_0: \sigma_1^2 \leq 24$ versus $H_a: \sigma_1^2 > 24$, the null hypothesis:

- ✓ **A) cannot be rejected.**
- X **B)** should neither be rejected nor fail to be rejected.
- X **C)** should be rejected.
- X **D)** cannot be tested using this sample information provided.

Explanation

The chi-square test is used to test hypotheses concerning a single population variance. Since this is a one-tailed test, the decision rule is to reject H_0 if $X^2 >$ the critical chi-square value at $\alpha = 0.05$ with $df = 24$.

$\chi^2_{n-1} = \frac{(n-1)s^2}{\sigma_0^2} = \frac{(24)25}{24} = 25.0$. The right-tail critical chi-square value is 36.415. Since $X^2 = 25 \leq 36.415$, H_0 cannot be rejected.

Question #41 of 82

Question ID: 438907

Consider the hypotheses structured as $H_0: \mu_1 \leq \$48$ versus $H_a: \mu_1 > \$48$. At a 5 percent level of significance, the null hypothesis:

- ☐ A) cannot be tested using the sample information provided.
- ☐ B) cannot be rejected.
- ☐ C) should neither be rejected nor fail to be rejected.
- ☒ D) should be rejected.

Explanation

A one-tailed t -test is appropriate. The decision rule is to reject H_0 if the computed t -statistic $>$ t -critical at $\alpha = 0.05$ with $df = 24$.

The computed value of the t -statistic = $\frac{\bar{x} - \mu_0}{s/\sqrt{n}} = \frac{50 - 48}{5/\sqrt{25}} = 2.0$, and t -critical = $t_{24} = 1.711$. Since $t >$ t -critical, H_0 should be rejected.

Question #42 of 82

Question ID: 438908

Using a 5 percent level of significance for a test of the null of $H_0: \sigma_1 = \sigma_2$ versus the alternative of $H_a: \sigma_1 \neq \sigma_2$, the null hypothesis:

- ☐ A) should neither be rejected nor fail to be rejected.
- ☐ B) should be rejected.
- ☐ C) cannot be tested using the sample information provided.
- ☒ D) cannot be rejected.

Explanation

The two-sided F -test is appropriate to test the equality of population variances. The decision rule is to reject H_0 if the computed test statistic, F , exceeds the critical F -value at $\alpha/2$. For the information provided, $F = s_1^2 / s_2^2 = 36/25 = 1.44$. At a 0.025 level of significance with $d_1 = 35$ and $d_2 = 24$, F -critical = 1.915. Since $F <$ F -critical ($1.44 < 1.915$), we fail to reject the null hypothesis.

Question #43 of 82

Question ID: 438819

A sample covariance for the common stock of the Earth Company and the S&P 500 is -9.50 . Which of the following statements regarding the estimated covariance of the two variables is *most* accurate?

- ☐ A) The two variables will have a strong tendency to move in opposite directions.
- ☐ B) The two variables will have a slight tendency to move together.
- ☒ C) The relationship between the two variables is not easily predicted by the calculated covariance.
- ☐ D) The relationship between the two variables is highly predictable given the calculated covariance.

Explanation

The actual value of the covariance for two variables is not very meaningful because its measurement is extremely sensitive to the scale of the two variables, ranging from negative to positive infinity. Covariance can, however be converted into the

correlation coefficient, which is more straightforward to interpret.

Question #44 of 82

Question ID: 438830

You observe the following annual returns on a stock portfolio for the last nine years: [5%, 7%, 20%, -4%, 2%, 5%, 13%, 18%, 4%]. Which measure of central tendency has the highest value?

- ☐ A) For these observations, the mean, median, and mode are equal.
- ☒ B) The mean.
- ☐ C) The mode.
- ☐ D) The median.

Explanation

Ordering the observations gives: [-4%, 2%, 4%, 5%, 5%, 7%, 13%, 18%, 20%]. The mode is the value that occurs most frequently; in this case, 5%. The median is the fifth ordered observation, 5%. The mean is: $(-4\% + 2\% + 4\% + 5\% + 5\% + 7\% + 13\% + 18\% + 20\%) / 9 = 7.8\%$.

Question #45 of 82

Question ID: 438848

The 95 percent confidence interval of the sample mean of the price earnings ratio for all traded stocks is 19 to 44. There are over 5,000 traded stocks and the sample size of this test is 100. Given that the expected value of the price earnings ratio is 31.5, the standard error of the ratio is *closest* to:

- ☐ A) 2.58.
- ☒ B) 6.38.
- ☐ C) 1.96.
- ☐ D) 12.50.

Explanation

The confidence interval is $31.5 \pm 1.96x$, where x is the standard error. If we take the upper bound, we know that $31.5 + 1.96x = 44$, or $1.96x = 12.5$. Hence, $x = 6.38$.

Question #46 of 82

Question ID: 438902

Abby Ness is an analyst for a firm that specializes in evaluating firms involved in mineral extraction. Ness believes that the earnings of copper extracting firms are more volatile than those of bauxite extraction firms. In order to test this, Ness examines the volatility of returns for 31 copper firms and 25 bauxite firms. The standard deviation of earnings for copper firms was \$2.69, while the standard deviation of earnings for bauxite firms was \$2.92. Ness's Null Hypothesis is $\sigma_1^2 = \sigma_2^2$. Based on the samples, can we reject the null hypothesis at a 95% confidence level using an F-statistic and why? Null is:

- ☐ A) rejected. The F-value exceeds the critical value by 0.71.
- ☐ B) rejected. The F-value exceeds the critical value by 0.849.
- ☐ C) not rejected. The critical value exceeds the F-value by 0.194.
- ☒ D) not rejected. The critical value exceeds the F-value by 0.71.

Explanation

$$F = s_1^2 / s_2^2 = \$2.92^2 / \$2.69^2 = 1.18$$

From an F table, the critical value with numerator df = 24 and denominator df = 30 is 1.89.

Question #47 of 82

Question ID: 438845

What is the *appropriate* test statistic for constructing confidence intervals for the population mean of a nonnormal distribution when the population variance is unknown and the sample size is large ($n \geq 30$)?

- ☐ A) The z-statistic at α with n degrees of freedom.
- ☐ B) The t -statistic at $\alpha/2$ with n degrees of freedom.
- ☒ C) The z-statistic or the t -statistic.
- ☐ D) The t -statistic at α with 29 degrees of freedom.

Explanation

When the sample size is large, and the central limit theorem can be relied upon to assure a sampling distribution that approaches normal, either the t -statistic or the z-statistic is acceptable for constructing confidence intervals for the population mean. However, the t -statistic will provide a more conservative range (wider) at a given level of significance.

Question #48 of 82

Question ID: 438901

You have collected monthly returns for a mutual fund and want to test the null hypothesis that the standard deviation exceeds the advertised standard deviation of 3.5 percent. The most appropriate test statistic is based on a:

- ☒ A) chi-square test.
- ☐ B) t-test.
- ☐ C) F-test.
- ☐ D) z-test.

Explanation

The chi-square test is used to test hypotheses about a single population variance.

Question #49 of 82

Question ID: 438877

A goal of an "innocent until proven guilty" justice system is to place a higher priority on:

- ☒ A) avoiding type I errors.
- ☐ B) the null hypothesis.
- ☐ C) avoiding type II errors.
- ☐ D) avoiding type III errors.

Explanation

In an "innocent until proven guilty" justice system, the null hypothesis is that the accused is innocent. The hypothesis can only be

rejected by evidence proving guilt beyond a reasonable doubt, favoring the avoidance of type I errors.

Question #50 of 82

Question ID: 438866

Maria Huffman is the Vice President of Human Resources for a large regional car rental company. Last year, she hired Graham Brickley as Manager of Employee Retention. Part of the compensation package was the chance to earn one of the following two bonuses: if Brickley can reduce turnover to less than 30%, he will receive a 25% bonus. If he can reduce turnover to less than 25%, he will receive a 50% bonus (using a significance level of 10%). The population of turnover rates is normally distributed. The population standard deviation of turnover rates is 1.5%. A recent sample of 100 branch offices resulted in an average turnover rate of 24.2%. Which of the following statements is *most* accurate?

- ☐ A) For the 50% bonus level, the critical value is -1.65 and Huffman should give Brickley a 50% bonus.
- ☐ B) Brickley should not receive either bonus.
- ☒ C) For the 50% bonus level, the test statistic is -5.33 and Huffman should give Brickley a 50% bonus.
- ☐ D) For the 25% bonus level, the test statistic is -10.66.

Explanation

Using the process of Hypothesis testing:

Step 1: State the Hypothesis. For 25% bonus level - $H_0: m \geq 30\%$ $H_a: m < 30\%$; For 50% bonus level - $H_0: m \geq 25\%$ $H_a: m < 25\%$.

Step 2: Select Appropriate Test Statistic. Here, we have a normally distributed population with a known variance (standard deviation is the square root of the variance) and a large sample size (greater than 30.) Thus, we will use the z-statistic.

Step 3: Specify the Level of Significance. $\alpha = 0.10$.

Step 4: State the Decision Rule. This is a one-tailed test. The critical value for this question will be the z-statistic that corresponds to an α of 0.10, or an area to the left of the mean of 40% (with 50% to the right of the mean). Using the z-table (normal table), we determine that the appropriate critical value = -1.28 (*Remember that we highly recommend that you have the "common" z-statistics memorized!*) Thus, we will reject the null hypothesis if the calculated test statistic is less than -1.28.

Step 5: Calculate sample (test) statistics. Z (for 50% bonus) = $(24.2 - 25) / (1.5 / \sqrt{100}) = -5.333$. Z (for 25% bonus) = $(24.2 - 30) / (1.5 / \sqrt{100}) = -38.67$.

Step 6: Make a decision. Reject the null hypothesis for both the 25% and 50% bonus level because the test statistic is less than the critical value. Thus, Huffman should give Soberg a 50% bonus.

The other statements are false. The critical value of -1.28 is based on the significance level, and is thus the same for both the 50% and 25% bonus levels.

Question #51 of 82

Question ID: 438820

The mean equity risk premium over a 40-year period is equal to 8.0 percent. The standard deviation of the sample is 12 percent. The standard error of the sample mean is *closest* to:

- X A) 1.26%.
- X B) 8.00%.
- X C) 0.30%.
- ✓ D) 1.90%.

Explanation

Note the size of the sample here is the number of years.

$$\sigma_{\bar{x}} = 12 / \sqrt{40} = 1.90\%$$

Question #52 of 82

Question ID: 438843

A nursery sells trees of different types and heights. Suppose that 75 pine trees are sold for planting at City Hall. These 75 trees average 60 inches in height with a standard deviation of 16 inches.

Using this information, construct a 95% confidence interval for the mean height of all trees in the nursery.

- X A) $0.8 \pm 1.96(16)$.
- ✓ B) $60 \pm 1.96(1.85)$.
- X C) $0.8 \pm 1.96(1.85)$.
- X D) $60 \pm 1.96(16)$.

Explanation

Because we know the population standard deviation, we use the z-statistic. A 95% confidence level is constructed by taking the population mean and adding and subtracting the product of the z-statistic reliability ($z_{\alpha/2}$) factor times the known standard deviation of the population divided by the square root of the sample size: $\bar{x} \pm z_{\alpha/2} \times (\sigma / n^{1/2}) = 60 \pm (1.96) \times (16 / 75^{1/2}) = 60 \pm (1.96) \times (16 / 8.6603) = 60 \pm (1.96) \times (1.85)$.

Question #53 of 82

Question ID: 438844

What is the *appropriate* test statistic for constructing confidence intervals for the population mean of a normal distribution when the population variance is unknown?

- X A) The z-statistic at α with n degrees of freedom.
- X B) The t-statistic at $\alpha/2$ with n degrees of freedom.
- X C) The z-statistic with $n - 1$ degrees of freedom.
- ✓ D) The t-statistic at $\alpha/2$ with $n - 1$ degrees of freedom.

Explanation

Use the t-statistic at $\alpha/2$ and $n - 1$ degrees of freedom when the population variance is unknown, regardless of sample size.

Question #54 of 82

Question ID: 438869

If the null hypothesis is innocence, then the statement "It is better that the guilty go free, than the innocent are punished" is an example of

preferring a:

- ☐ **A) two tailed test over a one tailed test.**
- ☐ **B) higher level of significance.**
- ☒ **C) type II error over a type I error.**
- ☐ **D) type I error over a type II error.**

Explanation

The statement shows a preference for not rejecting the null hypothesis when it is false (a type II error), over rejecting it when it is true (a type I error).

Question #55 of 82

Question ID: 438867

A bottler of iced tea wishes to ensure that an average of 16 ounces of tea is in each bottle. In order to analyze the accuracy of the bottling process, a random sample of 150 bottles is taken. Using a t -distributed test statistic of -1.09 and a 5% level of significance, the bottler should:

- ☐ **A) reject the null hypothesis and conclude that bottles do not contain an average of 16 ounces of tea.**
- ☐ **B) reject the null hypothesis and conclude that bottles contain an average 16 ounces of tea.**
- ☐ **C) not reject the null hypothesis and conclude that bottles do not contain an average of 16 ounces of tea.**
- ☒ **D) not reject the null hypothesis and conclude that bottles contain an average 16 ounces of tea.**

Explanation

$H_0: \mu = 16$; $H_a: \mu \neq 16$. Do not reject the null since $|t| = 1.09 < 1.96$ (critical value).

Question #56 of 82

Question ID: 438871

A Type I error:

- ☐ **A) fails to reject a true null hypothesis.**
- ☒ **B) rejects a true null hypothesis.**
- ☐ **C) fails to reject a false null hypothesis.**
- ☐ **D) rejects a false null hypothesis.**

Explanation

A Type I Error is defined as rejecting the null hypothesis when it is actually true. The probability of committing a Type I error is the significance level or alpha risk.

Question #57 of 82

Question ID: 438865

Which of the following statements about hypothesis testing is *least* accurate?

- ☐ **A) The null hypothesis is a statement about the value of a population parameter.**

- X **B)** A Type I error is rejecting the null hypothesis when it is actually true.
- ✓ **C)** If the alternative hypothesis is $H_a: \mu > \mu_0$, a two-tailed test is appropriate.
- X **D)** A Type II error is failing to reject a false null hypothesis.

Explanation

The hypotheses are always stated in terms of a population parameter. Type I and Type II are the two types of errors you can make - reject a null hypothesis that is true or fail to reject a null hypothesis that is false. The alternative may be one-sided (in which case a $>$ or $<$ sign is used) or two-sided (in which case a $\neq$ is used).

Question #58 of 82

Question ID: 438857

In the process of hypothesis testing, what is the proper order for these steps?

- X **A)** **Specify the level of significance. State the hypotheses. Make a decision. Collect the sample and calculate the sample statistics.**
- X **B)** State the hypotheses. Collect the sample and calculate the sample statistics. Make a decision. Specify the level of significance.
- X **C)** Collect the sample and calculate the sample statistics. State the hypotheses. Specify the level of significance. Make a decision.
- ✓ **D)** State the hypotheses. Specify the level of significance. Collect the sample and calculate the test statistics. Make a decision.

Explanation

The hypotheses must be established first. Then the test statistic is chosen and the level of significance is determined. Following these steps, the sample is collected, the test statistic is calculated, and the decision is made.

Question #59 of 82

Question ID: 438875

A return series with 250 observations has a sample mean of 10 percent and a standard deviation of 15 percent. The standard error of the sample mean is *closest* to:

- X **A)** 0.06.
- X **B)** 15.80.
- X **C)** 3.87.
- ✓ **D)** 0.95.

Explanation

The standard error of the sample mean is the standard deviation of the sample divided by the square root of the number of observations in the sample. In this case, $(15 / \sqrt{250}) = 0.95$.

Question #60 of 82

Question ID: 438836

The mean and variance of a sample average help define the distributional characteristics of a sample distribution and allow assumptions to be made about this distribution when the sample size is large. Which of the following choices represent the

standard deviation of a sample average?

- ☐ A) $\sigma(X)^2$
- ☐ B) $1/n$.
- ☒ C) $\sigma(X)/\sqrt{n}$.
- ☐ D) $\sigma(X)^2/n$.

Explanation

The central limit theorem states that for a population with a mean μ and a finite variance σ^2 , the sampling distribution of the sample mean of all possible samples of size n will be approximately normally distributed with a mean equal to μ and a variance equal to σ^2/n . Thus, the standard deviation of the sample average is equal to: $\sigma/\sqrt{n}$. This standard deviation measure is known as the standard error.

Question #61 of 82

Question ID: 438881

The mean monthly return for an equity portfolio over 60 months is 1.5%. The standard deviation is 3.0%. The value of the test statistic to test the hypothesis that mean monthly return is equal to zero is *closest* to:

- ☒ A) 3.87.
- ☐ B) 0.50.
- ☐ C) 2.19.
- ☐ D) 30.00

Explanation

$$z = \frac{1.5\% - 0.0\%}{\frac{3.0\%}{\sqrt{60}}} = 3.87.$$

Question #62 of 82

Question ID: 438855

Which of the following statements *least* describes the procedure for testing a hypothesis?

- ☐ A) **Select the level of significance, compute the test statistic, and make a decision.**
- ☐ B) Select the level of significance, formulate the decision rule, and make a decision.
- ☒ C) Compute the sample value of the test statistic, set up a rejection (critical) region, and make a decision.
- ☐ D) Develop a hypothesis, compute the test statistic, and make a decision.

Explanation

Depending upon the author there can be as many as seven steps in hypothesis testing which are:

1. Stating the hypotheses.
2. Identifying the test statistic and its probability distribution.
3. Specifying the significance level.
4. Stating the decision rule.
5. Collecting the data and performing the calculations.
6. Making the statistical decision.
7. Making the economic or investment decision.

Question #63 of 82

Question ID: 438880

Given the following hypothesis:

- The null hypothesis is $H_0 : \mu = 5$
- The alternative is $H_1 : \mu \neq 5$
- The mean of a sample of 17 is 7
- The population standard deviation is 2.0

What is the calculated z-statistic?

- ☐ A) 4.00.
- ☒ B) 4.12.
- ☐ C) 8.25.
- ☐ D) 8.00.

Explanation

The z-statistic is calculated by subtracting the hypothesized parameter from the parameter that has been estimated and dividing the difference by the standard error of the sample statistic. Here, the test statistic = (sample mean – hypothesized mean) / (population standard deviation / (sample size)^{1/2}) = $(\bar{X} - \mu) / (\sigma / n^{1/2}) = (7 - 5) / (2 / 17^{1/2}) = (2) / (2 / 4.1231) = 4.12$.

Question #64 of 82

Question ID: 438847

A random sample of 100 technology stocks earned an average of 10%. Assuming the distribution of equity returns is normal and the population standard deviation is 5%, the 95% confidence interval for the population mean is:

- ☐ A) 9.91% to 10.90%.
- ☐ B) 9.50% to 10.50%.
- ☒ C) 9.02% to 10.98%.
- ☐ D) 5.00% to 15.00%.

Explanation

$Z_{\alpha/2} = Z_{0.025} = 1.96$. So, $0.1 \pm 1.96(0.05 / 10) = 9.02\% \text{ to } 10.98\%$.

Question #65 of 82

Question ID: 438874

If the sample size is greater than 30 and population variance is unknown, the appropriate test for the sample mean is the:

- ☐ A) p-test or F-test.
- ☐ B) t-test.
- ☐ C) z-test.
- ☒ D) t-test or z-test.

Explanation

The central limit theorem makes it appropriate to use the z-test with an unknown variance if the sample size is large enough ($n \geq$

30), regardless of the distribution of the population. Since the t- and the z-distributions converge as sample size increases, either test is appropriate, although the t-test is a more conservative estimate.

Question #66 of 82

Question ID: 438833

A sample of paired points A and B is shown below. What is the covariance between the values of A and B?

Sample	A	B
1	1	2
2	4	5
3	9	7
4	11	10
5	14	12

X A) 82.20.

✓ B) 20.55.

X C) 7.80.

X D) 29.76.

Explanation

Sample	A	A-mean (A)	B	B-mean (B)	Product
1	1	-6.8	2	-5.2	35.36
2	4	-3.8	5	-2.2	8.36
3	9	1.2	7	-0.2	-0.24
4	11	3.2	10	2.8	8.96
5	14	6.2	12	4.8	29.76
mean	7.8		7.2	Sum	82.2

$$\text{Cov} = 82.20 / (5 - 1) = 20.55$$

Question #67 of 82

Question ID: 438856

The first step in the process of hypothesis testing is:

X A) the calculation of sample statistics.

X B) the collection of the sample.

X C) selecting the test statistic.

✓ D) to state the hypotheses.

Explanation

The researcher must state the hypotheses prior to the collection and analysis of the data. More importantly, it is necessary to

know the hypotheses before selecting the appropriate test statistic.

Question #68 of 82

Question ID: 438826

For the investments shown in the table below, what are the mean, median, and mode of the returns?

<i>Investment</i>	<i>Return (%)</i>
A	12
B	14
C	9
D	13
E	7
F	8
G	12

	<u>Mean</u>	<u>Median</u>	<u>Mode</u>
✓ A) 10.71%	12%	12%	12%
X B) 12.00%	12%	12%	12%
X C) 11.20%	12%	12%	12%
X D) 10.71%	9%	13%	13%

Explanation

The mean is the average return computed by summing the returns and dividing by the number of investments: $75 / 7 = 10.71\%$.

The median is the mid-point or central number of returns arranged from highest to lowest or lowest to highest. In this case: 7, 8, 9, 12, 12, 13, 14. The median return is 12%.

The mode is the return that occurs most frequently. In this case, 12% is also the mode.

Question #69 of 82

Question ID: 438850

A sample of size 25 is selected from a normal population. This sample has a mean of 15 and the population variance is 4.

Using this information, construct a 95% confidence interval for the population mean, μ .

- ✓ A) $15 \pm 1.96(0.4)$.
- X B) $15 \pm 1.96(2)$.
- X C) $15 \pm 1.96(4)$.
- X D) $15 \pm 1.96(0.8)$.

Explanation

Because we can compute the population standard deviation, we use the z-statistic. A 95% confidence level is constructed by taking the population mean and adding and subtracting the product of the z-statistic reliability ($z_{\alpha/2}$) factor times the known standard deviation of the population divided by the square root of the sample size (note that the population variance is given and

its positive square root is the standard deviation of the population): $x \pm z_{\alpha/2} \times (\sigma / n^{1/2}) = 15 \pm 1.96 \times (4^{1/2} / 25^{1/2}) = 15 \pm 1.96 \times (0.4)$.

Questions #70-71 of 82

Annual Returns on ABC Mutual Fund									
1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
11.0%	12.5%	8.0%	9.0%	13.0%	7.0%	15.0%	2.0%	-16.5%	11.0%

Question #70 of 82

Question ID: 438828

What is the arithmetic mean return for ABC Mutual Fund for the period 1991-2000?

Arithmetic Mean

- ☐ A) 28.2%
- ☐ B) 11%
- ☒ C) 7.2%
- ☐ D) 8.2%

Explanation

arithmetic mean = $(11 + 12.5 + 8 + 9 + 13 + 7 + 15 + 2 - 16.5 + 11) / 10 = 7.20$

Question #71 of 82

Question ID: 438829

Assuming a mean of 7.2%, what is the sample standard deviation of the returns for ABC Mutual Fund for the period 1991-2000?

- ☐ A) 7.8%.
- ☒ B) 9.1%.
- ☐ C) 10.2%.
- ☐ D) 9.8%.

Explanation

Standard deviation = $[\sum_i (x_i - \bar{X})^2 / (n - 1)]^{1/2}$

= $\sqrt{(744.10 / 9)} = \sqrt{(82.68)} = 9.1\%$

Question #72 of 82

Question ID: 438903

The sampling distribution of variances is described by which distribution?

- ☐ A) F-distribution.
- ☐ B) Student's t-distribution.
- ☐ C) Normal distribution.

✓ **D)** Chi-square distribution.

Explanation

The sampling distribution of variances is described by the chi-square distribution.

Question #73 of 82

Question ID: 438854

Which of the following is the correct sequence of events for testing a hypothesis?

- X **A)** State the hypothesis, select the level of significance, compute the test statistic, formulate the decision rule, and make a decision.
- ✓ **B)** State the hypothesis, select the level of significance, formulate the decision rule, compute the test statistic, and make a decision.
- X **C)** State the hypothesis, formulate the decision rule, select the level of significance, compute the test statistic, and make a decision.
- X **D)** State the hypothesis, formulate the decision rule, compute the test statistic, select the level of significance, and make a decision.

Explanation

Depending upon the author there can be as many as seven steps in hypothesis testing which are:

1. Stating the hypotheses.
 2. Identifying the test statistic and its probability distribution.
 3. Specifying the significance level.
 4. Stating the decision rule.
 5. Collecting the data and performing the calculations.
 6. Making the statistical decision.
 7. Making the economic or investment decision.
-

Question #74 of 82

Question ID: 438879

Which of the following statements is **FALSE**?

- X **A)** The probability of committing a Type I error is the significance level of the test.
- ✓ **B)** If a person is presumed innocent unless proven otherwise, finding a guilty person innocent is an example of a Type I error.
- X **C)** Failing to reject the null when it is false is an example of a Type II error.
- X **D)** A Type I error is rejecting the null when it is actually true.

Explanation

This is a Type II error. The null hypothesis is that the defendant is innocent which is not rejected when it should have been.

Question #75 of 82

Question ID: 438900

In order to test if Stock A is more volatile than Stock B, prices of both stocks are observed to construct the sample variance of the two

stocks. The appropriate test statistics to carry out the test is the:

- ☐ A) Chi-square test.
- ☐ B) Z test.
- ☒ C) F test.
- ☐ D) t test.

Explanation

The F test is used to test the differences of variance between two samples.

Question #76 of 82

Question ID: 438872

In a two-tailed test of a hypothesis concerning whether a population mean is zero, Jack Olson computes a t -statistic of 2.7 based on a sample of 20 observations where the distribution is normal. If a 5% significance level is chosen, Olson should:

- ☐ A) not make a conclusion pending additional observations.
- ☒ B) reject the null hypothesis and conclude that the population mean is significantly different from zero.
- ☐ C) reject the null hypothesis and conclude that the population mean is not significantly different from zero.
- ☐ D) fail to reject the null hypothesis that the population mean is not significantly different from zero.

Explanation

At a 5% significance level, the critical t -statistic using the Student's t -distribution table for a two-tailed test and 19 degrees of freedom (sample size of 20 less 1) is ± 2.093 (with a large sample size the critical z -statistic of 1.960 may be used). Because the critical t -statistic of 2.093 is to the left of the calculated t -statistic of 2.7, meaning that the calculated t -statistic is in the rejection range, we reject the null hypothesis and we conclude that the population mean is significantly different from zero.

Question #77 of 82

Question ID: 438831

The last three year-end returns for a stock are 5%, -2%, and 1%. Using an arithmetic mean, the sample standard deviation is *closest* to:

- ☐ A) 1.33%.
- ☐ B) 2.87%.
- ☐ C) 3.11%.
- ☒ D) 3.51%.

Explanation

The sample arithmetic mean is:

$$\bar{X} = (5 - 2 + 1) / 3 = 1.33\%$$

The sample standard deviation is:

$$s = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}} = \sqrt{\frac{(5\% - 1.33\%)^2 + (-2\% - 1.33\%)^2 + (1\% - 1.33\%)^2}{2}} = 3.51\%$$

Questions #78-82 of 82

XYZ Corp. Annual Stock Prices					
1995	1996	1997	1998	1999	2000
22%	5%	-7%	11%	2%	11%

Question #78 of 82

Question ID: 438838

What is the mean return for XYZ stock?

- ☐ A) 8.0%.
- ☐ B) -7.0%.
- ☒ C) 7.3%.
- ☐ D) 11.0%.

Explanation

$$\frac{[22\% + 5\% + (-7\%) + 11\% + 2\% + 11\%]}{6} = 7.3\%$$

Question #79 of 82

Question ID: 438839

Assuming that the distribution of XYZ stock returns is a population, what is the population variance (in %²)?

- ☐ A) 7.7.
- ☒ B) 80.2.
- ☐ C) 5.0.
- ☐ D) 6.8.

Explanation

The population variance, σ^2 , is found by taking the mean of all squared deviations from the mean.

$$\sigma^2 = \frac{[(22 - 7.3)^2 + (5 - 7.3)^2 + (-7 - 7.3)^2 + (11 - 7.3)^2 + (2 - 7.3)^2 + (11 - 7.3)^2]}{6} = 80.2(\%^2)$$

Question #80 of 82

Question ID: 438840

Assuming that the distribution of XYZ stock returns is a population, what is the population standard deviation?

- ☒ A) 8.96%.
- ☐ B) 5.02%.
- ☐ C) 46.22%.
- ☐ D) 6.84%.

Explanation

The population standard deviation, σ , is found by taking the square root of the population variance.

Question #81 of 82

Question ID: 438841

Assuming that the distribution of XYZ stock returns is a sample, what is the sample variance (in %²)?

- X A) 7.4.
- X B) 5.0.
- ✓ C) 96.3.
- X D) 72.4.

Explanation

The sample variance, s^2 , uses $n - 1$ in the denominator:

$$s^2 = \frac{(22 - 7.3)^2 + (5 - 7.3)^2 + (-7 - 7.3)^2 + (11 - 7.3)^2 + (2 - 7.3)^2 + (11 - 7.3)^2}{6 - 1} = 96.3(\%^2)$$

Question #82 of 82

Question ID: 438842

Assuming that the distribution of XYZ stock returns is a sample, what is the sample standard deviation?

- X A) 72.4%.
- X B) 7.4%.
- X C) 96.3%.
- ✓ D) 9.8%.

Explanation

The sample standard deviation is the square root of the sample variance:

$$s = \left\{ \frac{[(22 - 7.3)^2 + (5 - 7.3)^2 + (-7 - 7.3)^2 + (11 - 7.3)^2 + (2 - 7.3)^2 + (11 - 7.3)^2]}{6 - 1} \right\}^{1/2} = (96.3\%^2)^{1/2} = 9.8\%$$